

Control algebra for holonomic quantum computation with squeezed coherent states

Dennis Lucarelli

Department of Systems Science and Mathematics
Washington University, St. Louis, Missouri 63130

Abstract

A condition for determining the holonomy group associated to a principal bundle with connection is proven. Within the context of Holonomic Quantum Computation, this condition aids in determining the universality of the system. We apply the theory to the product bundle arising from the full two-qubit system of holonomic computation with squeezed coherent states and conclude that the control algebra generates the Lie algebra $u(4)$.

1 Introduction

Geometric phases have long been a source of fascination and insight into classical and quantum physical theories [1]. In recent years, they have proven to be useful in describing the dynamics and control of certain nonholonomic mechanical systems with symmetry. Inspired by the appearance of geometric phases in biology, engineers have sought to create motion in machines via cyclic variations in *shape space*. These endeavors [2] and the characterization of optimal trajectories [3, 4] remain active areas of research.

Most recently, the geometric phase has been realized as a way of constructing logic gates in a quantum computer [5]. Holonomic Quantum Computation (HQC) employs non-Abelian geometric phases (holonomies) for the purpose of quantum information processing. This paper is an elaboration of [6]. There it was reported that the *Optical Holonomic Computer* [7, 8, 9, 10, 11] is universal by showing that the holonomy group of the two-qubit interaction

model contains $SU(2) \times SU(2)$. These transformations act non-trivially on the standard computational basis and support, in particular, SWAP gates. Coupled with the ability to perform single qubit rotations, the claim of universality follows by appealing to standard results from quantum computation [12].

These notes and [6] completely solve the problem posed in the recent work of K. Fujii [9, 10, 11]. That is, for the two-qubit system, we explicitly construct a set of holonomic transformations that span the Lie algebra $u(4)$ and thus attain universality of the system. Note however that we do not carry out the program proposed in [11]. In Pachos-Chountasis [7] and Pachos-Zanardi [8], the universality of the model was shown by constructing a universal two-qubit logic gate. This $U(4)$ transformation was integrated directly by choosing a loop in a 2-submanifold of parameter space along which the connection components commute but with non-vanishing curvature. For these restricted loops, the non-Abelian nature of the path ordered integral may be avoided. However, that transformation was not an element of the holonomy group as calculated by the application of the Ambrose-Singer Theorem in [10].

In [6], we stated simple conditions for determining the holonomy group of a principal bundle with connection. These conditions are seen to be a corollary to Chow's theorem from control theory. The obvious relevance to HQC is that the holonomy group represents the set of controlled transformations attainable for manipulating quantum information.

In this section, we review some geometric constructions and provide some details regarding the corollary stated in [6]. This proof is not new. A similar proof can be found in [4] and, while not explicitly stated, all the necessary ingredients can be found in [13].

1.1 Principal and associated bundles

Let Q be a principal G -bundle defined by the free (right) Lie group action $\Phi : Q \times G \rightarrow Q$ and the canonical projection $Q \xrightarrow{\pi} Q/G \equiv M$. If U is a neighborhood of M , then Q is locally diffeomorphic to the product $U \times G$. A smooth map $\sigma : U \rightarrow Q$, such that $\pi \circ \sigma = \text{id}_M$ is called a local section over U .

Given any principal bundle $Q \xrightarrow{\pi} Q/G$ and an action of the Lie group G

on a vector space $\mathcal{V}$, we may define

$$\begin{aligned} \rho: Q \times \mathcal{V} &\rightarrow \mathcal{V} \\ (q, v) &\mapsto (qg, \rho^{-1}(g)v) \end{aligned} \quad (1)$$

where $q \in Q$, $v \in \mathcal{V}$, $g \in G$ and ρ is a representation of the group G . The *associated vector bundle* E_ρ with typical fiber $\mathcal{V}$ is the set of equivalence classes under this action

$$E_\rho = Q \times_\rho \mathcal{V} = (Q \times \mathcal{V})/G, \quad (2)$$

where points (q, v) and $(qg, \rho^{-1}(g)v)$ are identified. Let $[(q, v)]$ denote a representative of the equivalence class.

In the mathematical theory of HQC, there are *two* associated bundles of interest. First, the quantum code is realized as the $\mathbf{C}^n$ vector bundle associated to the pullback of the universal bundle by the classifying map [9, 10]. Second, the *adjoint bundle* defined by taking $\mathcal{V} = \mathfrak{g}$ (the Lie algebra of G) and ρ as the adjoint action of a Lie group on its Lie algebra is important in deriving conditions for universal computation as we prove in Section 1.3.

We now state an important correspondence concerning the associated adjoint bundle. First, recall that a Lie algebra valued differential form ω is Ad-equivariant if

$$\Phi_g^* \cdot \omega = \text{Ad}(g^{-1}) \cdot \omega \quad g \in G. \quad (3)$$

If ω is a Lie algebra valued *function* f on Q , then Ad-equivariance is written as $f(qg) = \text{Ad}(g^{-1})f(q)$, $g \in G$. The sections s of the adjoint bundle associated to a principal bundle Q are in one-to-one correspondence with Ad-equivariant functions on Q . To any Ad-equivariant function f we may define a local section s_f by $s_f(p) = [(q, f(q))]$, where q is any point in the fiber over p . This definition is independent of the choice of q because of Ad-equivariance. Conversely, given a local section $s: U \rightarrow Q \times_{\text{Ad}} \mathfrak{g}$, we define a map $f_s: \pi^{-1}(U) \rightarrow \mathfrak{g}$ such that $s(p) = [(p, f_s(p))]$ [14].

1.2 Connections and covariant derivatives

Recall that for any $\xi \in \mathfrak{g}$, the $\mathbf{R}$ -action $\Phi_{\exp(t\xi)} q$ defines a curve through $q \in Q$. The *infinitesimal generator* of the action corresponding to ξ is defined as the

vector field

$$\xi_q \equiv \left. \frac{d}{dt} \right|_{t=0} \Phi_{\exp(t\xi)} q. \quad (4)$$

The vertical subspace $V_q Q \subset T_q Q$ is defined to be the subspace of $T_q Q$ that is tangent to the fiber over $p = \pi(q)$, G_p . By the previous definition, we have the identification $V_q Q \cong \mathfrak{g}$.

A connection on a principal G -bundle $\mathcal{A}$ on Q is an Ad-equivariant Lie algebra-valued one-form $\mathcal{A} : TQ \rightarrow \mathfrak{g}$ such that $A(\xi_q) = \xi$. The horizontal space $H_q Q$ is the linear space $H_q Q \equiv \{X \in T_q Q | \mathcal{A}_q(X) = 0\}$. These definitions provide the splitting

$$T_q Q = V_q Q \oplus H_q Q \quad (5)$$

of the tangent vectors into horizontal and vertical components. Note that

$$V_q Q = \text{Ker } T_q \pi, \quad \text{and} \quad H_q Q = \text{Ker } \mathcal{A}_q. \quad (6)$$

The map $T_q \pi : H_q \rightarrow T_{\pi(q)} M$ is an isomorphism from the horizontal space to the tangent space of the base space. Thus a curve $q(t) \in Q$ defines a curve in the base by specifying a tangent vector at each point $\pi(q(t)) = p(t)$. The properties of the connection and the uniqueness theory of ODE's provide the reverse procedure of reconstructing a unique curve in the horizontal space given a curve in the base called the *horizontal lift* [15]. Denote the horizontal lift of the vector $X \in T_p M$ where $p = \pi(q)$ by X_q^h and $h X_q$ as the horizontal component of a vector in $T_q Q$.

Let σ be a local section over a neighborhood U of M that contains a curve γ . Then the horizontal lift $\tilde{\gamma}$ can be expressed as $\tilde{\gamma}(t) = \sigma(\tilde{\gamma}(t))g(t)$, where $g(t) = g(\gamma(t)) \in G$. The local connection one-form A is defined as the pullback $A \equiv \sigma^*(\mathcal{A})$. The local connection may be written in terms of group elements by the equation

$$\frac{dg(t)}{dt} = -A g(t). \quad (7)$$

Choosing the initial condition, $g(0) = \text{id}_G$, the solution may be obtained by the path ordered integral

$$g(\gamma(t)) = \mathbf{P} \exp \int_{\gamma} -A. \quad (8)$$

Given a connection on a principal bundle, we may define the covariant derivative of sections of associated bundles. Let $E_{\mathfrak{g}} = Q \times_{\text{Ad}} \mathfrak{g}$ be the adjoint bundle associated to Q as defined above. Let γ be a base curve with tangent vector X at $\gamma(0)$. If $\{\mathcal{T}_\alpha\}$ is a basis of $\mathfrak{g}$, then a section s of $E_{\mathfrak{g}}$ may be obtained by forming $s = [(\sigma(p), \mathcal{T}(p))]$, where $\mathcal{T}(p) = f^\alpha(p)\mathcal{T}_\alpha$ and σ is a local section over a neighborhood U of p . The covariant derivative $D_X s$ of s along $X \in TM$ is defined by [15]

$$D_X s \equiv [(\tilde{\gamma}(0), \left. \frac{d}{dt} \text{Ad}_{g^{-1}(t)} \mathcal{T}(t) \right|_{t=0})] \quad (9)$$

The previous definitions imply,

$$D_X s = [(\sigma(0)g(0), \left. \frac{d}{dt} \text{Ad}_{g^{-1}(t)} \mathcal{T}(t) \right|_{t=0})] \quad (10)$$

$$= [(\sigma(0), \left. \text{Ad}_g \frac{d}{dt} \text{Ad}_{g^{-1}(t)} \mathcal{T}(t) \right|_{t=0})] \quad (11)$$

$$= [(\sigma(0), \left. \frac{d\mathcal{T}}{dt} + [A(X), \mathcal{T}(t)] \right|_{t=0})]. \quad (12)$$

Covariant derivatives of sections of the adjoint bundle provide a way of computing Lie derivatives of Ad-equivariant functions on the total space by the correspondence

$$D_X s_f = X^h f \quad (13)$$

where s_f is the section of the adjoint bundle corresponding to the Ad-equivariant function f [13].

1.3 Infinitesimal holonomy

The horizontal lift of any closed curve in the base space with initial point p maps the fiber over p to itself and thus corresponds to the group translation $qg, q \in \pi^{-1}(p)$ and $g \in G$. In this way, we can associate to closed curves in the base space elements of the structure group G . The set of all such group elements forms a subgroup of G called the *holonomy group*. With the restriction to sufficiently small loops homotopic to zero, we may define the *infinitesimal* holonomy group. The infinitesimal holonomy group is in general a subgroup of the full holonomy group, however for analytic connections over connected base manifolds, these notions coincide [13]

The holonomy group is determined from the curvature. The *curvature* $\mathcal{F}$ associated to $\mathcal{A}$ is by definition the Lie algebra-valued two-form on Q such that

$$\mathcal{F}(X_1, X_2) = \mathbf{d}\mathcal{A}(h X_1, h X_2) \quad (14)$$

where $\mathbf{d}$ denotes exterior differentiation. Using the identity

$$\mathbf{d}\mathcal{A}(X_1, X_2) = X_1\mathcal{A}(X_2) - X_2\mathcal{A}(X_1) - \mathcal{A}([X_1, X_2]) \quad (15)$$

and the fact that the connection form evaluates to zero on horizontal vectors, we get the relation $\mathcal{F}(X_1, X_2) = -\mathcal{A}([h X_1, h X_2])$. Roughly, the curvature measures the non-commutativity in the total space of horizontally lifted vector fields defined in the base space. Note that $\mathcal{F}$ is Ad-equivariant since $\mathcal{A}$ is Ad-equivariant and pullback and exterior differentiation commute.

The Lie algebra corresponding to the infinitesimal holonomy group is spanned by the $\mathfrak{g}$ -valued Ad-equivariant function f given by

$$f = X_{i_k}^h X_{i_{k-1}}^h \dots X_{i_3}^h \mathcal{F}(X_{i_2}^h, X_{i_1}^h) \quad (16)$$

where $X_i \in TM$ and $2 \leq k < \infty$ [16]. By the correspondence (13), the pullback through a local section σ of this function may be written in terms of covariant derivatives as

$$D_{X_{i_k}} D_{X_{i_{k-1}}} \dots D_{X_{i_3}} F(X_{i_2}, X_{i_1}) \quad (17)$$

where F is the local form of the curvature defined with respect to a local section $F \equiv \sigma^*(\mathcal{F})$.

2 HQC with squeezed coherent states

In this section, we revisit the mathematical framework of Optical Holonomic Computation. See [7, 8, 9, 10] for the many details omitted here. We refer to this setup as *HQC with squeezed coherent states* since the properties of interest in quantum computation, e.g. universality, entanglement, etc., rely only on the squeezing and displacing operators and not directly on the degenerate Hamiltonian used to encode the qubits. Therefore, the mathematics describing this computing scheme will be shared by any quantum system whose states can be displaced and squeezed. See for example the remarks concerning trapped ions in [8].

Let $\mathcal{H}_i = \{|\alpha_i\rangle, \alpha \in \mathbf{N}\}$ be a Fock space with number operator n_i . The Hamiltonian $H^i = n_i(n_i - 1)$ encodes the i -th qubit in the degenerate subspace $\{|0_i\rangle, |1_i\rangle\}$. Single mode squeezing and displacing operators control the single qubit

$$S^i(\mu_i) = \exp(\mu_i a_i^{\dagger 2} - \bar{\mu}_i a_i^2) \quad D^i(\lambda_i) = \exp(\lambda_i a_i^\dagger - \bar{\lambda}_i a_i) \quad (18)$$

where $\mu_i, \lambda_i \in \mathbf{C}$ and a_i and $a_i^\dagger$ are the annihilation and creation operators respectively. To obtain controllable interactions among product states we employ the Hamiltonian $H^{12} = n_1(n_1 - 1) + n_2(n_2 - 1)$ with degenerate subspace $\{|00\rangle, |01\rangle, |10\rangle, |11\rangle\}$ and two mode squeezing and displacing operators

$$M(\zeta) = \exp(\zeta a_1^\dagger a_2^\dagger - \bar{\zeta} a_1 a_2) \quad N(\xi) = \exp(\xi a_1^\dagger a_2 - \bar{\xi} a_1 a_2^\dagger) \quad (19)$$

where $\xi, \zeta \in \mathbf{C}$. Set $\mathcal{U}_2 = D(\lambda)S(\mu)$ and $\mathcal{U}_4 = N(\xi)M(\zeta)$.

The geometry of the $U(n)$ holonomy problem [9, 10] is briefly described as follows. Define the manifolds

$$\text{St}_n(\mathcal{H}) = \{V = (v_1, \dots, v_n) \in \mathcal{H} \times \dots \times \mathcal{H} \mid V^\dagger V = \text{Id}_{n \times n}\} \quad (20)$$

$$\text{Gr}_n(\mathcal{H}) = \{X \in B(\mathcal{H}) \mid X^2 = X, X^\dagger = X, \text{tr} X = n\}. \quad (21)$$

where $B(\mathcal{H})$ denotes the set of bounded linear operators on $\mathcal{H}$. These manifolds are known as the Stiefel and Grassmann manifolds respectively. They form a principal bundle with the (right) $U(n)$ action on $\text{St}_n(\mathcal{H})$ and the projection $\pi : \text{St}_n(\mathcal{H}) \rightarrow \text{Gr}_n(\mathcal{H})$ given by $\pi(V) = VV^\dagger$. Denote this $U(n)$ -bundle by P_n . Let $M \subset \mathbf{C}^2$ be a parameter space and define the classifying map $\Pi_n : M \rightarrow \text{Gr}_n(\mathcal{H})$ given by

$$\Pi_n(\eta) \equiv \mathcal{U}_n(\eta) \left(\sum_{j=1}^n v_j v_j^\dagger \right) \mathcal{U}_n^\dagger(\eta) \quad (22)$$

Now form the pullback bundle $Q_n = \Pi_n^* P_n$. The quantum code is then realized as the $\mathbf{C}^n$ -vector bundle associated to Q_n [9, 10, 15]. To ease the notation, we consider only the manifolds P_n and Q_n . So then, the full two-qubit system consisting of single qubit rotations on each qubit and interactions among them is modelled by the pullback bundles corresponding to the product bundle

$$\begin{array}{ccccc} \text{St}_2(\mathcal{H}_1) & \times & \text{St}_4(\mathcal{H}_1 \otimes \mathcal{H}_2) & \times & \text{St}_2(\mathcal{H}_2) \\ \downarrow & & \downarrow & & \downarrow \\ \text{Gr}_2(\mathcal{H}_1) & \times & (\text{Gr}_4)^{\text{int}}(\mathcal{H}_1 \otimes \mathcal{H}_2) & \times & \text{Gr}_2(\mathcal{H}_2) \end{array} \quad (23)$$

The local connection one-form is given by the Wilczek-Zee connection [17] with matrix elements

$$A_{\eta_i}^{\bar{\varrho}\varrho} \equiv \langle \bar{\varrho} | \mathcal{U}^\dagger(\eta) \frac{\partial}{\partial \eta_i} \mathcal{U}(\eta) | \varrho \rangle. \quad (24)$$

The connection components are listed in Appendix A with respect to the labelling of parameters $\lambda_1 = x_1 + iy_1$, $\mu_1 = r_1 e^{i\theta_1}$, $\lambda_2 = x_2 + iy_2$, $\mu_2 = r_2 e^{i\theta_2}$, $\zeta = r_3 e^{i\theta_3}$ and $\xi = r_4 e^{i\theta_4}$.

2.1 Reduction

The local curvature components are listed in Appendix B. The holonomy group of the two-qubit interaction model is $SU(2) \times SU(2) \times U(1)$ [6]. These transformations leave invariant the subspaces $\{|00\rangle, |11\rangle\}$ and $\{|01\rangle, |10\rangle\}$ in $\mathcal{H}_1 \otimes \mathcal{H}_2$. By the reduction theorem for connections [13], the connection is reducible to a $su(2) \times su(2) \times u(1)$ -valued connection and we may reduce the total space to $\text{St}_{2,4}(\mathcal{H}_1 \otimes \mathcal{H}_2)$. To determine the reduced base manifold, we form the quotient

$$\frac{U(4)}{SU(2) \times SU(2) \times U(1)} \cong \frac{SU(4)}{SU(2) \times SU(2)} \cong \text{SGr}_{2,4} \equiv (\text{Gr}_{2,4})^{\text{int}}. \quad (25)$$

In a similar manner, we can reduce the bundles $\text{St}_2(\mathcal{H}_i) \mapsto \text{Gr}_2(\mathcal{H}_i)$ corresponding to the single qubit rotations. The $U(2)$ holonomies act in the product space $\mathcal{H}_1 \otimes \mathcal{H}_2$ as $U(2) \otimes \mathbf{1}$ and $\mathbf{1} \otimes U(2)$. The bundles reduce to $\text{St}_{2,4}(\mathcal{H}_1 \otimes \mathcal{H}_2) \mapsto \text{Gr}_{2,4}(\mathcal{H}_1 \otimes \mathcal{H}_2)$. For the full two-qubit system, we have the reduced product bundle

$$\begin{array}{ccccc} \text{St}_{2,4}(\mathcal{H}_1 \otimes \mathcal{H}_2) & \times & \text{St}_{2,4}(\mathcal{H}_1 \otimes \mathcal{H}_2) & \times & \text{St}_{2,4}(\mathcal{H}_1 \otimes \mathcal{H}_2) \\ \downarrow & & \downarrow & & \downarrow \\ \text{Gr}_{2,4}(\mathcal{H}_1 \otimes \mathcal{H}_2) & \times & (\text{Gr}_{2,4})^{\text{int}}(\mathcal{H}_1 \otimes \mathcal{H}_2) & \times & \text{Gr}_{2,4}(\mathcal{H}_1 \otimes \mathcal{H}_2) \end{array} \quad (26)$$

This structure is similar to the product bundle for the generic two-qubit system proposed by Pachos [18].

2.2 Control algebra

The standard argument to prove universality of a quantum system is to demonstrate the ability to perform all single qubit rotations on each qubit

and a non-trivial two-qubit transformation. The implicit argument is then that the *control algebra* formed by these matrices and their Lie brackets generates the Lie algebra $su(n)$ or $u(n)$. In this section, we carry out this argument for HQC with squeezed coherent states. The matrices we refer to can be found in Appendices B and C.

The holonomy algebra for each single qubit system is $u(2)$.

$$u(2) \otimes \mathbf{1} : \text{span}_{\mathbf{R}}\{F_{x_1 y_1} \otimes \mathbf{1}, F_{r_1 \theta_1} \otimes \mathbf{1}, F_{y_1 r_1} \otimes \mathbf{1}, F_{x_1 r_1} \otimes \mathbf{1}\} \quad (27)$$

$$\mathbf{1} \otimes u(2) : \text{span}_{\mathbf{R}}\{\mathbf{1} \otimes F_{x_2 y_2}, \mathbf{1} \otimes F_{r_4 \theta_4}, \mathbf{1} \otimes F_{y_2 r_4}, \mathbf{1} \otimes F_{x_2 r_4}\} \quad (28)$$

The holonomy algebra for two qubit interactions system is $su(2) \times su(2) \times u(1)$.

$$\text{span}_{\mathbf{R}}\{F_{r_2 r_3}, F_{r_2 \theta_2}, F_{r_2 \theta_3}, F_{r_3 \theta_3}, D_{\frac{\partial}{\partial r_2}} F_{r_2 \theta_2}, D_{\frac{\partial}{\partial \theta_2}} F_{r_2 \theta_2}, D_{\frac{\partial}{\partial \theta_2}} D_{\frac{\partial}{\partial \theta_2}} F_{r_2 \theta_2}\} \quad (29)$$

From these matrices, we choose the subset linearly independent over the reals

$$\mathcal{C}_1 = \{F_{r_1 \theta_1} \otimes \mathbf{1}, F_{y_1 r_1} \otimes \mathbf{1}, F_{x_1 r_1} \otimes \mathbf{1}\} \quad (30)$$

$$\mathcal{C}_2 = \{\mathbf{1} \otimes F_{x_2 y_2}, \mathbf{1} \otimes F_{y_2 r_4}, \mathbf{1} \otimes F_{x_2 r_4}\} \quad (31)$$

$$\mathcal{C}_{12} = \{F_{r_2 r_3}, F_{r_2 \theta_2}, F_{r_2 \theta_3}, D_{\frac{\partial}{\partial r_2}} F_{r_2 \theta_2}, D_{\frac{\partial}{\partial \theta_2}} D_{\frac{\partial}{\partial \theta_2}} F_{r_2 \theta_2}\} \quad (32)$$

Additional independent matrices can be obtained by taking Lie brackets from the set $\{\mathcal{C}_1 \cup \mathcal{C}_2 \cup \mathcal{C}_{12}\}$. In principle, these transformations can be constructed in the usual infinitesimal way. For example, to construct $[F_{x_1 r_1} \otimes \mathbf{1}, F_{r_2 r_3}]$ traverse a small loop in the plane (x_1, r_1) , then traverse a loop in the plane (r_2, r_3) , then traverse the loop backwards in (x_1, r_1) , and finally traverse the loop backwards in (r_2, r_3) . One choice is given by

$$\begin{aligned} \mathcal{C}_{12}^2 = \{ & [F_{x_1 r_1} \otimes \mathbf{1}, F_{r_2 r_3}], [\mathbf{1} \otimes F_{x_2 r_4}, F_{r_2 r_3}], [F_{y_1 r_1} \otimes \mathbf{1}, F_{r_2 \theta_3}], \\ & [\mathbf{1} \otimes F_{y_2 r_4}, F_{r_2 \theta_3}], [\mathbf{1} \otimes F_{y_2 r_4}, [F_{x_1 r_1} \otimes \mathbf{1}, F_{r_2 r_3}]] \} \end{aligned} \quad (33)$$

Note that these matrices are independent with each other and the set $\{\mathcal{C}_1 \cup \mathcal{C}_2 \cup \mathcal{C}_{12}\}$ over the reals.

The 16 matrices $\mathcal{C} = \{\mathcal{C}_1 \cup \mathcal{C}_2 \cup \mathcal{C}_{12} \cup \mathcal{C}_{12}^2\}$ span the Lie algebra $u(4)$. Thus, at least in principle, we have obtained complete controllability over the two-qubit system.

Appendix

A Connection components

A.1 Single qubit

$$\begin{aligned}
 A_{x_1} &= \begin{bmatrix} -iy_1 & -(\cosh 2r_1 - e^{i\theta_1} \sinh 2r_1) \\ \cosh 2r_1 - e^{-i\theta_1} \sinh 2r_1 & -iy_1 \end{bmatrix} \\
 A_{y_1} &= \begin{bmatrix} ix_1 & i(\cosh 2r_1 + e^{i\theta_1} \sinh 2r_1) \\ i(\cosh 2r_1 + e^{-i\theta_1} \sinh 2r_1) & ix_1 \end{bmatrix} \\
 A_{\theta_1} &= \begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix} \frac{i}{4} (\cosh 4r_1 - 1) \quad A_{r_1} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}
 \end{aligned}$$

$$\begin{aligned}
 A_{x_2} &= \begin{bmatrix} -iy_2 & -(\cosh 2r_4 - e^{i\theta_4} \sinh 2r_4) \\ \cosh 2r_4 - e^{-i\theta_4} \sinh 2r_4 & -iy_2 \end{bmatrix} \\
 A_{y_2} &= \begin{bmatrix} ix_2 & i(\cosh 2r_2 + e^{i\theta_4} \sinh 2r_4) \\ i(\cosh 2r_4 + e^{-i\theta_4} \sinh 2r_4) & ix_2 \end{bmatrix} \\
 A_{\theta_4} &= \begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix} \frac{i}{4} (\cosh 4r_4 - 1) \quad A_{r_4} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}
 \end{aligned}$$

A.2 Two-qubit

$$A_{r_2} = \begin{bmatrix} 0 & 0 & 0 & -e^{-i\theta_2} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ e^{i\theta_2} & 0 & 0 & 0 \end{bmatrix} \quad A_{r_3} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & -e^{-i\theta_3} & 0 \\ 0 & e^{i\theta_3} & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} (2 \cosh^2 r_2 - 1)$$

$$\begin{aligned}
 A_{\theta_2} &= \begin{bmatrix} 0 & 0 & 0 & e^{-i\theta_2} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ e^{i\theta_2} & 0 & 0 & 0 \end{bmatrix} \frac{i}{2} \sinh 2r_2 + \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \end{bmatrix} \frac{i}{2} (\cosh 2r_2 - 1) \\
 A_{\theta_3} &= \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & e^{-i\theta_3} & 0 \\ 0 & e^{i\theta_3} & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \frac{i}{2} \cosh 2r_2 \sin 2r_3 + \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} i \sin^2 r_3
 \end{aligned}$$

B Holonomy algebra

B.1 $u(2) \otimes \mathbf{1}$

$$\begin{aligned}
F_{x_1 y_1} &= \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} 4i & F_{r_1 \theta_1} &= \begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix} i \sinh 4r_1 \\
F_{y_1 r_1} &= \begin{bmatrix} 0 & -1 \\ 0 & 0 \end{bmatrix} 2i(e^{i\theta_1} \cosh 2r_1 + \sinh 2r_1) \\
&+ \begin{bmatrix} 0 & 0 \\ -1 & 0 \end{bmatrix} 2i(e^{-i\theta_1} \cosh 2r_1 + \sinh 2r_1) \\
F_{x_1 r_1} &= \begin{bmatrix} 0 & -1 \\ 0 & 0 \end{bmatrix} 2(e^{i\theta_1} \cosh 2r_1 - \sinh 2r_1) \\
&+ \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} 2(e^{-i\theta_1} \cosh 2r_1 - \sinh 2r_1) \\
\\
F_{x_1 y_1} \otimes \mathbf{1} &= \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} 4i & F_{r_1 \theta_1} \otimes \mathbf{1} &= \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 3 \end{bmatrix} i \sinh 4r_1 \\
\\
F_{y_1 r_1} \otimes \mathbf{1} &= \begin{bmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} 2i(e^{i\theta_1} \cosh 2r_1 + \sinh 2r_1) \\
&+ \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{bmatrix} 2i(e^{-i\theta_1} \cosh 2r_1 + \sinh 2r_1) \\
\\
F_{x_1 r_1} \otimes \mathbf{1} &= \begin{bmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} 2(e^{i\theta_1} \cosh 2r_1 - \sinh 2r_1) \\
&+ \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix} 2(e^{-i\theta_1} \cosh 2r_1 - \sinh 2r_1)
\end{aligned}$$

B.2 $\mathbf{1} \otimes u(2)$

$$F_{x_2 y_2} = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} 4i \quad F_{r_4 \theta_4} = \begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix} i \sinh 4r_4$$

$$F_{y_2 r_4} = \begin{bmatrix} 0 & -1 \\ 0 & 0 \end{bmatrix} 2i(e^{i\theta_4} \cosh 2r_4 + \sinh 2r_4) \\ + \begin{bmatrix} 0 & 0 \\ -1 & 0 \end{bmatrix} 2i(e^{-i\theta_4} \cosh 2r_4 + \sinh 2r_4)$$

$$F_{x_2 r_4} = \begin{bmatrix} 0 & -1 \\ 0 & 0 \end{bmatrix} 2i(e^{i\theta_4} \cosh 2r_4 - \sinh 2r_4) \\ + \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} 2i(e^{-i\theta_4} \cosh 2r_4 - \sinh 2r_4)$$

$$\mathbf{1} \otimes F_{x_2 y_2} = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} 4i \quad \mathbf{1} \otimes F_{r_4 \theta_4} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 3 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 3 \end{bmatrix} i \sinh 4r_4$$

$$\mathbf{1} \otimes F_{y_2 r_4} = \begin{bmatrix} 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \end{bmatrix} 2i(e^{i\theta_4} \cosh 2r_4 + \sinh 2r_4) \\ + \begin{bmatrix} 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \end{bmatrix} 2i(e^{-i\theta_4} \cosh 2r_4 + \sinh 2r_4) \\ \mathbf{1} \otimes F_{x_2 r_4} = \begin{bmatrix} 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \end{bmatrix} 2i(e^{i\theta_4} \cosh 2r_4 - \sinh 2r_4) \\ + \begin{bmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix} 2i(e^{-i\theta_4} \cosh 2r_4 - \sinh 2r_4)$$

B.3 $su(2) \times su(2) \times u(1)$

$$\begin{aligned}
F_{r_2 r_3} &= \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & -e^{-i\theta_3} & 0 \\ 0 & e^{i\theta_3} & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} 2 \sinh 2r_2 & F_{r_2 \theta_2} &= \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 2 \end{bmatrix} 2i \sinh 2r_2 \\
F_{r_2 \theta_3} &= \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & e^{-i\theta_3} & 0 \\ 0 & e^{i\theta_3} & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} i \sin 2r_3 \sinh 2r_2 & F_{r_3 \theta_3} &= \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} i \sin 2r_3 \sinh^2 2r_2, \\
D_{\frac{\partial}{\partial \theta_2}} F_{r_2 \theta_2} &= \begin{bmatrix} 0 & 0 & 0 & -e^{-i\theta_2} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ e^{i\theta_2} & 0 & 0 & 0 \end{bmatrix} 2 \sinh^2 2r_2 \\
D_{\frac{\partial}{\partial r_2}} F_{r_2 \theta_2} &= \begin{bmatrix} 0 & 0 & 0 & -e^{-i\theta_2} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -e^{i\theta_2} & 0 & 0 & 0 \end{bmatrix} 4i \sinh 2r_2 + \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 2 \end{bmatrix} 4i \cosh 2r_2, \\
D_{\frac{\partial}{\partial \theta_2}} D_{\frac{\partial}{\partial \theta_2}} F_{r_2 \theta_2} &= \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix} 2i \sinh^3 r_2 + \begin{bmatrix} 0 & 0 & 0 & e^{-i\theta_2} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ e^{i\theta_2} & 0 & 0 & 0 \end{bmatrix} 2i \sinh^2 2r_2 \cosh 2r_2
\end{aligned}$$

C Higher order brackets

$$\begin{aligned}
[F_{x_1 r_1} \otimes \mathbf{1}, F_{r_2 r_3}] &= \begin{bmatrix} 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} 4e^{i(\theta_1+\theta_3)} \sinh 2r_2 (\cosh 2r_1 - e^{-i\theta_1} \sinh 2r_1) \\
&+ \begin{bmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \end{bmatrix} 4e^{-i(\theta_1+\theta_3)} \sinh 2r_2 (\cosh 2r_1 - e^{i\theta_1} \sinh 2r_1) \\
[\mathbf{1} \otimes F_{x_2 r_4}, F_{r_2 r_3}] &= \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} 4e^{-i(\theta_3-\theta_4)} \sinh 2r_2 (\cosh 2r_4 - e^{-i\theta_4} \sinh 2r_4) \\
&+ \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix} 4e^{i(\theta_3-\theta_4)} \sinh 2r_2 (\cosh 2r_4 - e^{i\theta_4} \sinh 2r_4) \\
[F_{y_1 r_1} \otimes \mathbf{1}, F_{r_2 \theta_3}] &= \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \end{bmatrix} 2e^{i(\theta_1+\theta_3)} \sin 2r_3 \sinh 2r_2 (\cosh 2r_1 + e^{-i\theta_1} \sinh 2r_1) \\
&+ \begin{bmatrix} 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} 2e^{-i(\theta_1+\theta_3)} \sin 2r_3 \sinh 2r_2 (\cosh 2r_1 + e^{i\theta_1} \sinh 2r_1) \\
[\mathbf{1} \otimes F_{y_2 r_4}, F_{r_2 \theta_3}] &= \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} 2e^{-i(\theta_3-\theta_4)} \sin 2r_3 \sinh 2r_2 (\cosh 2r_4 + e^{-i\theta_4} \sinh 2r_4) \\
&+ \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix} 2e^{i(\theta_3-\theta_4)} \sin 2r_3 \sinh 2r_2 (\cosh 2r_4 + e^{i\theta_4} \sinh 2r_4)
\end{aligned}$$

$$\left[\mathbf{1} \otimes F_{y_2 r_4}, [F_{x_1 r_1} \otimes \mathbf{1}, F_{r_2 r_3}] \right] = \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix} \delta$$

$$\delta = 16i \sinh 2r_2 \left(-\sinh 2r_1 (\cos(\theta_3 - \theta_4) \cosh 2r_4 + \cos \theta_3 \sinh 2r_4) \right. \\ \left. + \cosh 2r_1 (\cos(\theta_1 + \theta_3 - \theta_4) \cosh 2r_4 + \cos(\theta_1 + \theta_3) \sinh 2r_4) \right)$$

Acknowledgements : I would like to thank SML and MKM without whose cooperation this work would not have been possible.

References

- [1] A. Shapere and F. Wilczek eds. 1989 *Geometric Phases in Physics* (Singapore: World Scientific)
- [2] S. Kelly and R. Murray 1995 J. Robotic Syst. **12** 417
- [3] W. Koon and J. Marsden 1997 SIAM J. Cont. Opt. **35** 901
- [4] R. Montgomery 2002 *A Tour of Subriemannian Geometries, Their Geodesics, and Applications*, (Providence: American Mathematical Society)
- [5] P. Zanardi and M. Rasetti 1999 Phys. Lett. A **264** 94 quant-ph/9904011
- [6] D. Lucarelli *Chow's theorem and universal holonomic quantum computation* quant-ph/0111078
- [7] J. Pachos and S. Chountasis 2000 Phys. Rev. A **62** 052318 quant-ph/9912093
- [8] J. Pachos and P. Zanardi 2001 Int. J. Mod. Phys. B **15** 1257 quant-ph/0007110
- [9] K. Fujii 2000 J. Math. Phys. **41** 4406 quant-ph/9910069
- [10] K. Fujii 2001 Rep. Math. Phys. **48** 75 quant-ph/0004102

- [11] K. Fujii 2001 *Mathematical foundations of holonomic quantum computer II* quant-ph/0101102
- [12] M. Nielsen and I. Chuang, 2000 *Quantum Computation and Quantum Information* (New York: Cambridge University Press)
- [13] S. Kobayashi and K. Nomizu 1969 *Foundations of Differential Geometry* vol. 1 (New York: Wiley Interscience)
- [14] K. Marathe and G. Martucci, 1991 *The Mathematical Foundations of Gauge Theories* (Amsterdam: North Holland)
- [15] M. Nakahara 1990 *Geometry, Topology, and Physics* (Bristol: IOP Publishing Ltd.)
- [16] H. Ozeki 1956 Nagoya Math. J. **10** 106.
- [17] F. Wilczek and A. Zee 1984 Phys. Rev. Lett. **52** 2111
- [18] J. Pachos 2000 *Quantum computation by geometrical means* quant-ph/0003150